

**USE OF AERIAL PHOTOGRAPHY TO ASSESS
CHANGES IN THE DISTRIBUTION OF ELKHORN CORAL
IN THE U.S. VIRGIN ISLANDS**

PREPARED FOR:

U.S. Geological Survey
Biological Resources Division
USGS Caribbean Field Station
St. John, U.S. Virgin Islands

PREPARED BY:

Jean-Pierre Bacle
island resources
FOUNDATION
St. Thomas, U.S. Virgin Islands

September 2002

TABLE OF CONTENTS

| | PAGE |
|---------------------------------------|------|
| OBJECTIVES | 1 |
| BACKGROUND | 1 |
| METHODOLOGY | 1 |
| LIMITATIONS | 3 |
| RESULTS | 4 |
| REMARKS & RECOMMENDATIONS | 8 |
| ACKNOWLEDGEMENTS | 9 |
| REFERENCES | 10 |
| APPENDIX #1: Figure A and B | |
| APPENDIX #2: Aerial Photo Compilation | |
| APPENDIX #3: Figures #1 to # 14 | |

LIST OF TABLES

| | | |
|----------|---|---|
| Table #1 | Selected imagery for the five study sites | 2 |
| Table #2 | Hawksnest Bay, <i>A. palmata</i> coverage | 5 |
| Table #3 | Windswept, <i>A. palmata</i> coverage | 6 |
| Table #4 | Haulover Bay, <i>A. palmata</i> coverage | 7 |
| Table #5 | Newfound Bay, <i>A. palmata</i> coverage | 7 |
| Table #6 | Buck Island, <i>A. palmata</i> coverage | 8 |

USE OF AERIAL PHOTOGRAPHY TO ASSESS CHANGES IN THE DISTRIBUTION OF ELKHORN CORAL IN THE U.S. VIRGIN ISLANDS

OBJECTIVES

The main purpose of this project is to examine the feasibility of using conventional aerial photographs as a tool to document the historical and current distribution of *Acropora palmata* (Elkhorn) coral colonies. The study also attempts to document the health of *A. palmata* wherever possible. Five pre-selected “Pilot Areas” in the U.S. Virgin Islands were selected for this study. They include: Hawksnest Bay, Windswept, Haulover Bay, Newfound Bay and Buck Island (see Figures A and B in Appendix #1).

BACKGROUND

Coral reef systems at depths of 0–5m around St. John, St. Thomas and St. Croix were once dominated by large colonies of branching coral *A. palmata* commonly known as Elkhorn coral. Elkhorn coral grows rapidly (5-10cm/yr), and has a complex morphology which provide conditions conducive to support a highly diverse fish community and habitat for many other reef organisms. The rapid decline of *A. palmata* has primarily been attributed to necrosis associated with white band disease and physical destruction from tropical storms and hurricanes. The combination of these factors, white band disease and storm damage, have contributed to reductions in live coverage of *A. palmata* by up to 80% in many locations of the USVI.

METHODOLOGY

To date, one of the most cost effective technologies for mapping shallow water benthic habitats is through the use of conventional aerial photo interpretation assisted with GIS based image analysis. Aerial photographs were used to develop the Benthic Habitats of the Florida Key digital data atlas (EMRI, 1998) and just recently, a similar effort was performed for the U.S. Virgin Islands and Puerto Rico as part of the National Ocean Service’s continuing effort to document coastal resources (Kendall, *et al.*, 2001).

The following methodology was adapted for the specific objectives of the project. The work was broken out into five main tasks summarized as follows.

1) **Aerial Photo Selection Process**

A search to gather all information regarding the availability and location of historical and recent aerial photographs from the National Ocean Service was conducted at the onset of this study. The effort resulted in a detailed compilation of available aerial color photographs arranged according to location and date flown (see Appendix 2).

From the compiled list, selective aerial photographs for the five study sites were chosen. The selection process was based on two main criteria: optimum scale and quality of imagery. The largest scales available ranged between 1:12,000 and 1:20,000. The image quality selection process considered the amount of cloud cover, shade, sun glint, turbidity and sea state.

To evaluate changes in the *A. palmata* community through time, historical images from the 1970s were selected to compare with the most up to date coverage (1999). Coverage from the 1980s was also chosen to examine if any trend occurred between these two dates. The process resulted in a final selection as shown in the table below.

Table #1: Selected imagery for the five study sites

| Study Sites | 1971 1:20,000 | 1974 1:12,000 | 1977 1:20,000 | 1983 1:15,000 | 1999 1:48,000 |
|--------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| Hawksnest Bay | | • | | • | • |
| Windswept | | • | | • | • |
| Haulover Bay | • | | | • | • |
| Newfound Bay | • | | | • | • |
| Buck Island | | | • | | • |

2) **Digital Imagery**

Aerial photos from the five study areas (1971 to 1983) were scanned with a high-resolution scanner at 1,200 dots per inch (DPI) and orthorectified. The 1999 imagery was previously scanned at 500 DPI. The georeference digital imagery output formed the basis from which the GIS Arc View 3.1 image analysis was applied.

3) **Aerial Photo Interpretation**

Conventional aerial photo interpretation with the use of a stereoscope was applied to both hard copy color imagery and diapositives (color transparencies) placed on a light table. This initial exercise allowed for an overview and establish rough delineation of fringing and barrier reef systems and preliminary assessment of *A. palmata* distribution.

4) Field Verification

Reconnaissance site visits were undertaken to validate aerial photo observations from the most current coverage (1999). Sites were visually evaluated by snorkeling and/or from the boat in shallow and clear water. Because of time and budget limitation, Buck Island was excluded.

5) Image Analysis and Mapping

Analysis and mapping was performed with the use of Arc View 3.1 that included an image analysis extension. The image analysis extension allowed for the use of a host of tools which could enhance the interpretation's accuracy and confidence level. Such tools included:

- Adjusting the brightness and contrast of the image
- Choosing band combinations
- Enhancing image display
- Applying custom histogram stretches to obtain specific visual results
- Sharpening image appearance
- Smoothing image appearance
- Using edge detection function

All mapping was conducted by using "heads-up" on-screen digitizing. Digitizing scale was done at approximately 1:2,500 to 1:3,000 for the sites. The digitized product was subsequently reduced to fit an 8.5 by 11 hardcopy presentation. A total of 16 maps were produced for the five sites (see Figures #1 to #14 in Appendix 3).

LIMITATIONS

Aerial photo interpretation with the support of GIS image analysis technology offers great potential to identify and delineate shallow water benthic communities. In this study however, we found that the interpretation's accuracy and confidence levels were ultimately linked to the scale and quality of the imagery.

Scale Limitation

Small scale imagery where ground objects appear small will have less potential to yield details. In this study, the largest scale available was 1:12,000 for only two of the selected pilot areas (see table #1). At this scale, one millimeter measured on the photographs translated to 12 meters at the ground level. Assuming that mature *A. palmata* colonies could only reach 1m to 2 m in diameter, visual recognition was deemed unreliable at this scale with identifying individual heads, however stands greater than 9 square meters

could likely be identified. Increasing the scale with the GIS “zoom-in” function did little to help identify individual stands. At this small scale, the physical dimension of the *A. palmata* feature did not yield a clear signature reading (distinctive shape, pattern and color tone recognition).

Variability in scale from 1:12,000 to 1:48,000 also introduced an element of inconsistency in the mapping (digitizing) process. Since “on-screen” digitizing was done at approximately 1:2,500 for all sites, greater delineation accuracy was accomplished on the larger scale photos than the 1:48,000 coverage.

Aerial Photo Quality

Other factors limiting the accuracy of the interpretation and mapping included shadow fall, sun glint, cloud cover, turbidity and sea state. These factors were carefully considered in the process of choosing the best imagery, however, in some cases were unavoidable. For example, the 1983 coverage had significant “shadow fall” along the coastline that hindered the interpretation (see Figures #5, #8 and #11 in Appendix 3). It appears that the 1983 coverage was flown in late afternoon. That factor combined with the rugged coastline and slope aspect projected shadow falls in many sites.

Digital image quality also varied due to different scanning resolution. The ready available 1999 (1:48,000) coverage was scanned at a resolution of 500 dots per inch (DPI) while all other coverage selected for this study was scanned at 1,200 DPI resolution. The 1999 coverage with its lower scanning resolution and with its significantly smaller scale resulted in a lower image quality (fuzzier image) considered inadequate for this kind of study application.

RESULTS

Despite the severe limiting factors mentioned above, the study provided some useful information as to the distribution and relative abundance of *A. palmata*. It also provided some indication of historical trends (see Tables #1 to #5 and Figures #1 to #14 in Appendix 3).

With aerial photo interpretation combined with image analysis, it was possible to delineate the reef boundary where *A. palmata* generally occurs. These zones included the upper fore reef, the breaker zone and the reef flat.

In using the image analysis tools and the “zoom-in” function, it was possible to delineate areas densely covered with aggregate stands of *A. palmata* colonies. Distribution and abundance were estimated but with limited degree of accuracy. In most of these areas the *A. palmata* cover was intermixed with other coral species such as *Millepora sp.*, *Montastrea sp* and other fragmented coral and rubble. For this reason, dense cover of *A.*

palmata stands was mapped under a 60-80% cover class. Shallow reef areas covered with scattered *A. palmata* stands were mapped under the <10% class cover.

Discerning health conditions of *A. palmata* colonies such as White Band Disease (WBD) was not feasible at any scale. Identifying “standing dead” stands would be possible at a larger photo scale because shape and structure is still preserved. It was possible however to outline extensive zones that have been transformed into rubble fields due to past storms and/or areas of ongoing high energy waves constantly reworking the coral rubble. These areas usually have a high reflectance value that can be detected on the imagery.

The 5 pilot areas are described below with observations related to abundances displayed in a tabular fashion. Corresponding figures #1 to #14 are located in Appendix 3.

(1) **Hawksnest Bay**

Hawksnest Bay is a semi-enclosed bay and is somewhat sheltered from the normal condition of easterly wave and swell for most of the year. During the winter months, however, the northern swell is common and contributes much turbulence and wave action on the upper fore reef and beach zone.

Hawksnest has a series of shallow reefs that occupy the inner bay area. Three are narrow, elongated and extend perpendicular from the sandy beach. The fourth is situated along the eastern edge of the bay, near Gibney’s beach. Total reef area is approximately 9,945m².

Image analysis indicates that in 1974, 11.5% of the reef areas were densely covered with *A. palmata* (60% to 80% cover). Analysis of the 1983 photos reveal a slight reduction to 10.4% and down to 5.6% by 1999 (see Table #2 and Figures #1, #2 and #3 in Appendix 3).

Table #2: Hawksnest Bay, *A. palmata* coverage

| | 1974 1:12,000 | | 1983 1:15,000 | | 1999 1:48,000 | |
|--------------------------------------|--|--|-------------------------------------|--|-------------------------------------|-------------------------------------|
| | < 10% cover | 60% - 80% cover | <10 % cover | 60% - 80% cover | <10% cover | 60% - 80% cover |
| Reef Area 1 3,748 m ² | 734m ² | 362m ² | 35m ² | 426m ² | | 385m ² |
| Reef Area 2 3,653 m ² | 694m ² | 536m ² | 75m ² | 448m ² | 534m ² | 270m ² |
| Reef Area 3 1,121 m ² | | 130m ² | 51m ² | | 108m ² | |
| Reef Area 4 1,423 m ² | | 111m ² | 75m ² | 160m ² | 132m ² | |
| Total 9,945 m² | 1,428 m² (14.3%) | 1,139 m² (11.5%) | 236 m² (2.3%) | 1,034 m² (10.4%) | 774 m² (7.7%) | 566 m² (5.6%) |

(2) Windswept

Windswept is located immediately east of Trunk Bay. The area can be characterized as an exposed rocky headland fronted by a large fringing reef system. The area is protected from easterly currents by Mary's Point. During the winter months the northern swell causes breaking waves on the reef.

The fringing reef can be subdivided into four areas, the larger ones separated by distinctive sand channels. The total reef system covers about 39,735 m². From the 1974 imagery, it is estimated that dense *A. palmata* areas (60% to 80%) occupied 9.5% of the total reef area. In 1983, the dense cover was reduced to 3% and increased back to 6.4% by 1999 (see Table #4 and Figures #4, #5 and #6). The significant decrease in 1983 is partially due to poor image quality as a result of shadow fall obstructing the interpretation (see Figure #5).

Table #3: Windswept, *A. palmata* coverage

| | 1974 1:12,000 | | 1983 1:15,000 | | 1999 1:48,000 | |
|---------------------------------------|---|---------------------------------------|--|---------------------------------------|--|---------------------------------------|
| | <10% cover | 60%-80% cover | <10%cover | 60%-80% cover | <10%cover | 60%-80% cover |
| Reef Area 1 11,550 m ² | 3,928m ² | 213m ² | 3,842m ² | N/A | 3,524m ² | 571m ² |
| Reef Area 2 19,185 m ² | 5,253m ² | 2,144m ² | 3,219m ² | 821m ² | 4,410m ² | 1,078m ² |
| Reef Area 3 3,500 m ² | 1,273m ² | 178m ² | 1,389m ² | N/A | 1,110m ² | 152qm ² |
| Reef Area 4 5,500 m ² | 1,103m ² | 1,264m ² | 1,273m ² | 360m ² | 842m ² | 751m ² |
| Total 39,735 m² | 11,557 m² (29.0%) | 3,799 m² (9.5%) | 9,723 m² (24.4%) | 1,181 m² (2.9%) | 9,886 m² (24.8%) | 2,552 m² (6.4%) |

(3) Haulover Bay

Haulover Bay is a large, partially exposed bay of greater depth. The image analysis was limited to the western side of the bay which is within the park boundary. The eastern portion of the bay is predominantly subtidal bedrock.

The western reef system is breached in two by a narrow sand channel. The combined reef area cover is 15,650 m². Analysis of the 1971 imagery estimate extensive *A. palmata* thickets (60% to 80%) covered about 26% of the reef area. The recent 1999 imagery revealed that dense cover has completely disappeared (see Table #4 and Figures 7, 8 and 9). Area #1 in Figure #8 is not shown due to a wrong frame selection.

Table #4: Haulover Bay, *A. palmata* cover

| | 1971 1:20,000 | | 1983 1:15,000 | | 1999 1:48,000 | |
|---------------------------------------|---------------------------------------|--------------------------------------|--|---------------------------------------|---------------------------------------|---------------|
| | <10% cover | 60%-80% cover | <10% cover | 60%-80% cover | <10% cover | 60%-80% cover |
| Reef Area 1 10,696 m ² | 8,013m ² | 2,683m ² | Incomplete Image | Incomplete Image | 1980m | 0 |
| Reef Area 2 4,954 m ² | 3,583m ² | 1,371m ² | 3,842m ² | 1,112m ² | 0 | 0 |
| Total 15,650 m² | 11,596 m² (74%) | 4,054 m² (26%) | 3,842 m² (24.5%) | 1,112 m² (7.1%) | 1980 m² (12.6%) | 0 |

(4) **Newfound Bay**

Located in the east end of St John along the north shore, Newfound Bay benefits from the sheltering effect of a partially closed bay mouth barrier reef system. Since 1971, the exposed reef system has suffered significant decline in live coral including *A. palmata*. In 1971 it is estimated that 16% of this reef was densely covered (60% to 80%). By 1983, densely covered areas were reduced to a little over 9% of the reef area. Imagery from 1999 show no aggregate stands, only sparse occurrences (see Table #5 and Figures #10, #11 and #12).

Figures #12a and #12b represent examples of selective image analysis tools that were used in the interpretation.

Despite the obvious wipeout, field reconnaissance during 2002 revealed numerous individual stands of young colonies sprouting all along the reef crest and upper fore reef. Most of the stands were estimated at 30cm in diameter.

Table #5: Newfound Bay, *A. palmata* cover

| | 1971 1:20,000 | | 1983 1:15,000 | | 1999 1:48,000 | |
|---------------------------------------|---|--|---|---------------------------------------|-------------------------------------|---------------|
| | <10% cover | 60%-80% cover | <10% cover | 60%-80% cover | <10% cover | 60%-80% cover |
| Reef Area 1 8,258 m ² | 6,906m ² | 1,352m ² | 7,332m ² | 926m ² | 155m ² | none |
| Reef Area 2 13,635 m ² | 10,453m ² | 2,182m ² | 12,583m ² | 1,052m ² | none | none |
| Total 21,893 m² | 17,359 m² (79.2%) | 3,534 m² (16.1%) | 19,915 m² (90.9%) | 1,978 m² (9.0%) | 155 m² (0.7%) | none |

(5) **Buck Island**

Buck Island is located 2 km north of Teague Bay, St Croix. The study area includes the barrier reef that wraps around the eastern tip of the island and is situated approximately 200-250 m from the shoreline. The location and structure of this barrier reef forms an arc that protects Buck Island from the dominant easterly wave attack.

The study area is limited to the shallow portion of the barrier reef, namely the reef crest, upper fore reef and adjoining back reef. It is segmented into seven reef areas forming a total of 113,974 m². Aerial photo analysis indicates that in 1977, 30.6% of the study area was densely covered with *A. palmata* thickets (60% to 80% cover). Recent 1999 imagery reveals a dramatic decline in that only 9.2% of the study area has very sparse (<10%) occurrences of *A. palmata* cover (see Table #6 and Figures #13 and #14).

No photo coverage was available from the 1980s.

Table #6: Buck Island, *A. palmata* cover

| | 1977 (1:20,000) | 1999 (1:48,000) |
|------------------------------------|--|---------------------------------------|
| | 60%-80% cover | <10% cover |
| Reef Area 1 1,220m ² | 315m ² | 231m ² |
| Reef Area 2 8,595m ² | 1,301m ² | 999m ² |
| Reef Area 3 3,450m ² | 1,014m ² | 885m ² |
| Reef Area 4 7,330m ² | 1,780m ² | 1,397m ² |
| Reef Area 5 7,486m ² | 2,875m ² | 2,445m ² |
| Reef Area 6 77,430m ² | 25,300m ² | 3,583m ² |
| Reef Area 7 8,463m ² | 2,367m ² | 923m ² |
| Total: 113,974m² | 34,952m² (30.6%) | 10,464m² (9.2%) |

REMARKS & RECOMMENDATIONS

Aerial photo interpretation has proven to be a useful tool to define and delineate benthic habitat. Limits in the accuracy of the interpretation do exist and are usually attributed to the scale and quality of the image as well as resolution lost due to the scanning process. It's also important to note that the success in carrying out the aerial photo interpretation exercise depends in large measure on prior training and experience of the interpreter in the discipline relevant to the problem in question. Thus, it's reasonable to expect that no two interpreters will produce the exact same results.

This study concludes that identifying individual *A. palmata* colonies is not possible from the aerial photos available even with the aid of a GIS “zoom-in” function and analysis tools. The aerial photo scales are considered too small for this type of application. Furthermore, variability in scale from historical coverage to present is considerable and creates inherent inconsistencies in the interpretation process.

It is possible however to interpret and delineate extensive, densely aggregated stands of *A. palmata* with a moderate level of accuracy. A classification cover of (60% to 80% cover) was set to better reflect densely aggregate stands of *A. palmata* cover. It is important to note that such areas also include other coral species in the mix such as *Millepora sp* and *Montastrea sp* as well as variable quantities of fragmented and dead coral. As a result, this study should be considered more as a qualitative assessment rather than a quantitative one.

To better meet the objectives of such a study, acquisition of large scale photography is a prerequisite. Minimum aerial photo scale should be at least 1:500, meaning that one millimeter (mm) measured on the aerial photo translates to 0.5m on the ground. Form, structure and signature tone of individual *A. palmata* stands could be identified at that scale and perhaps its health status also.

The cost benefit for such a study remains to be determined. The very first priority should focus on designing a list of specifications tailored specifically for this kind of application. Priority considerations should be on low altitude large scale imagery and optimum image type. Conventional color is adequate, however other image types should be reviewed such as the Compact Airborne Spectrographic Imager (CASI) system. It was recently reported that using this special digital sensor system on a plane flying at 250 meters above sea level helped diagnose more comprehensively the health status of reefs (Raloff, 2001).

ACKNOWLEDGEMENTS

The following persons were instrumental in providing technical support and guidance:

Dr. Caroline Rogers, U.S. Geological Survey

Dr. Barry Devine, Center for Data Conservation, University of the U.S. Virgin Islands

Tim Battista, NOAA National Ocean Service

Dr. Marc Monaco, NOAA National Ocean Service

Matt Kendall, NOAA National Ocean Service

Ken Buja, NOAA National Ocean Service

Lita Katz, NOAA National Ocean Service

Marilyn King, NOAA National Ocean Service

REFERENCES

- Anderson, M., Lund, H., Gladfelter, E., and M. Davis. 1986. Ecological community type maps and biological descriptions for the Buck Island Reef National Monument and proposed marine sites in the British Virgin Islands. Biosphere Reserve Report Number 4, National Park Service, Dept. of Interior. 235pp.
- Bacle, J.P. 1995. Mapping coastal habitat features Great Pond Bay, St. Croix. U.S. Virgin Islands. *Photo Interpretation*. 4:264-268.
- Beets, J., L. Leeward, and E.S. Zullo. 1986. Marine community descriptions and maps of the bays within the Virgin Islands National Park/Biosphere Reserve. Biosphere Reserve Research Report Number 2, National Park Service, Dept. of Interior. 118pp.
- Florida Fish and Wildlife Conservation Commission, Florida Marine Research Institute (FMRI) and National Ocean and Atmospheric Administration. 1998. Benthic Habitats of the Florida Keys. FMRI Technical Report No. TR-4. 52pp.
- Kendall, M.S., C.R. Kruer, K.R. Buja, J.D. Christensen, M. Finkbeiner, and M.E. Monaco. 2001. NOAA Technical Memorandum NOS NCCOS CCMA 152. Methods used to map the benthic habitats of Puerto Rico and the U.S. Virgin Islands. 45pp.
- Raloff, J., 2001. Aircraft spies on health of coral reefs. *Science News*, Vol.160, No.10.
- Rogers, C., 2000. Is *Acropora palmata* (Elkhorn coral) making a comeback in the Virgin Islands? *Reef Encounter* 27, July.

Appendix 2: Aerial Photo Compilation

| Roll No. | Photo | Scale | Date | Map Index No. | Area Covered | Quality | incomplete | glint | shade | clouds | turbidity |
|----------|-------|-------|------------|---------------|--------------|---------|------------|-------|-------|--------|-----------|
| 100-720 | 846 | 1:30k | 11/15/1971 | PR3-4 | Windswept | A | | | | | |
| 100-720 | 847 | 1:30k | 11/15/1971 | PR3-4 | Hawksnest | A | | | | | |
| 100-720 | 847 | 1:30k | 11/15/1971 | PR3-4 | Windswept | A | | | | | |
| 100-720 | 848 | 1:30k | 11/15/1971 | PR3-4 | Hawksnest | A | | | | | |
| 100-720 | 848 | 1:30k | 11/15/1971 | PR3-4 | Windswept | C | | X | | | |
| 100-720 | 889 | 1:30k | 11/15/1971 | PR3-4 | Hawksnest | B | | | | X | |
| 100-720 | 889 | 1:30k | 11/15/1971 | PR3-4 | Windswept | A | | | | | |
| 100-720 | 890 | 1:30k | 11/15/1971 | PR3-4 | Hawksnest | B | | | | X | |
| 100-720 | 890 | 1:30k | 11/15/1971 | PR3-4 | Windswept | A | | | | | |
| 100-720 | 891 | 1:30k | 11/15/1971 | PR3-4 | Windswept | A | | | | | |
| 100-720 | 893 | 1:30k | 11/15/1971 | PR3-4 | Haulover | B | | X | | | |
| 100-720 | 893 | 1:30k | 11/15/1971 | PR3-4 | Newfound | B | | X | | | |
| 100-720 | 894 | 1:30k | 11/15/1971 | PR3-4 | Haulover | B | | X | | | |
| 100-720 | 894 | 1:30k | 11/15/1971 | PR3-4 | Newfound | B | | X | | | |
| 100-720 | 895 | 1:30k | 11/15/1971 | PR3-4 | Haulover | A | | X | | | |
| 100-720 | 895 | 1:30k | 11/15/1971 | PR3-4 | Newfound | A | | X | | | |
| 100-720 | 897 | 1:30k | 11/15/1971 | PR3-4 | Windswept | B | | | | X | |
| 100-720 | 898 | 1:30k | 11/15/1971 | PR3-4 | Hawksnest | A | | | | | |
| 100-720 | 898 | 1:30k | 11/15/1971 | PR3-4 | Windswept | A | | | | X | |
| 100-720 | 899 | 1:30k | 11/15/1971 | PR3-4 | Hawksnest | B | | X | | | |
| 100-720 | 899 | 1:30k | 11/15/1971 | PR3-4 | Windswept | B | | X | | | |
| 100-720 | 914 | 1:30k | 11/15/1971 | PR3-4 | Haulover | C | X | | X | X | |
| 100-720 | 915 | 1:30k | 11/15/1971 | PR3-4 | Haulover | C | X | X | X | | |
| 100-720 | 915 | 1:30k | 11/15/1971 | PR3-4 | Newfound | A | | | | X | |
| 100-721 | 966 | 1:20k | 11/15/1971 | PR3-3 | Hawksnest | C | X | X | | | |
| 100-721 | 966 | 1:20k | 11/15/1971 | PR3-3 | Windswept | C | | X | | X | |
| 100-721 | 967 | 1:20k | 11/15/1971 | PR3-3 | Hawksnest | C | X | X | | | |
| 100-721 | 967 | 1:20k | 11/15/1971 | PR3-3 | Windswept | C | | X | | | |
| 100-721 | 1000 | 1:30k | 11/20/1971 | PR3-4 | None | X | | | | | |

Use of Aerial Photography to Assess Changes in Distribution of Elkhorn Coral in the USVI

| Roll No. | Photo | Scale | Date | Map Index No. | Area Covered | Quality | incomplete | glint | shade | clouds | turbidity |
|----------|-------|-------|------------|---------------|--------------|---------|------------|-------|-------|--------|-----------|
| 100-721 | 1001 | 1:30k | 11/20/1971 | PR3-4 | Haulover | C | X | | X | | |
| 100-721 | 1001 | 1:30k | 11/20/1971 | PR3-4 | Newfound | A | | | | | |
| 100-721 | 1002 | 1:30k | 11/20/1971 | PR3-4 | Haulover | A | | | | | |
| 100-721 | 1002 | 1:30k | 11/20/1971 | PR3-4 | Newfound | B | | X | | | |
| 100-721 | 1003 | 1:30k | 11/20/1971 | PR3-4 | Haulover | A | | | | | |
| 100-721 | 1003 | 1:30k | 11/20/1971 | PR3-4 | Newfound | B | | X | X | | |
| 100-721 | 1031 | 1:30k | 11/20/1971 | PR3-4 | Hawksnest | A | | | | | |
| 100-721 | 1031 | 1:30k | 11/20/1971 | PR3-4 | Windswept | A | | | | | |
| 100-721 | 1032 | 1:30k | 11/20/1971 | PR3-4 | Hawksnest | A | | | | | |
| 100-721 | 1032 | 1:30k | 11/20/1971 | PR3-4 | Windswept | A | | | | | |
| 100-721 | 1033 | 1:30k | 11/20/1971 | PR3-4 | Windswept | A | | | | | |
| 100-721 | 1035 | 1:30k | 11/20/1971 | PR3-4 | Haulover | A | | | | | |
| 100-721 | 1035 | 1:30k | 11/20/1971 | PR3-4 | Newfound | B | | X | | | |
| 100-721 | 1036 | 1:30k | 11/20/1971 | PR3-4 | Haulover | A | | | | | |
| 100-721 | 1036 | 1:30k | 11/20/1971 | PR3-4 | Newfound | A | | X | | | |
| 100-721 | 1037 | 1:30k | 11/20/1971 | PR3-4 | None | X | | | | | |
| 100-721 | 1045 | 1:20k | 11/20/1971 | PR3-3 | Haulover | A | | | X | | |
| 100-721 | 1046 | 1:20k | 11/20/1971 | PR3-3 | Haulover | A | | X | | | |
| 100-721 | 1052 | 1:20k | 11/20/1971 | PR3-3 | Haulover | A | | X | | | |
| 100-721 | 1052 | 1:20k | 11/20/1971 | PR3-3 | Newfound | C | | X | | | |
| 100-721 | 1053 | 1:20k | 11/20/1971 | PR3-3 | Haulover | A | | | | | |
| 100-721 | 1053 | 1:20k | 11/20/1971 | PR3-3 | Newfound | A | | X | | | |
| 100-721 | 1054 | 1:20k | 11/20/1971 | PR3-3 | Newfound | A | | | | | |
| 100-722 | 1123 | 1:30k | 11/20/1971 | PR3-3 | Buck | B | X | | | | |
| 100-722 | 1124 | 1:30k | 11/20/1971 | PR3-3 | Buck | A | X | | | | |
| 100-722 | 1125 | 1:30k | 11/20/1971 | PR3-3 | Buck | B | X | | | | |
| 100-823 | 7014 | 1:12k | 2/11/1974 | PR3-6 | Hawksnest | B | X | | | | |
| 100-823 | 7015 | 1:12k | 2/11/1974 | PR3-6 | Hawksnest | A | | | | | |
| 100-823 | 7016 | 1:12k | 2/11/1974 | PR3-6 | Hawksnest | A | | | | | |
| 100-824 | 7103 | 1:12k | 2/12/1974 | PR3-6 | Hawksnest | A | | | | | |


Use of Aerial Photography to Assess Changes in Distribution of Elkhorn Coral in the USVI

| Roll No. | Photo | Scale | Date | Map Index No. | Area Covered | Quality | incomplete | glint | shade | clouds | turbidity |
|----------|-------|-------|------------|---------------|--------------|---------|------------|-------|-------|--------|-----------|
| 100-824 | 7104 | 1:12k | 2/12/1974 | PR3-6 | Hawksnest | A | | | | | |
| 100-824 | 7111 | 1:12k | 2/12/1974 | PR3-6 | Windswept | A | | | | | |
| 100-824 | 7112 | 1:12k | 2/12/1974 | PR3-6 | Windswept | A | | | | | |
| 100-824 | 7113 | 1:12k | 2/12/1974 | PR3-6 | Hawksnest | A | | | | | |
| 100-824 | 7113 | 1:12k | 2/12/1974 | PR3-6 | Windswept | B | X | | | | |
| 100-824 | 7114 | 1:12k | 2/12/1974 | PR3-6 | Hawksnest | A | | | | | |
| 100-824 | 7115 | 1:12k | 2/12/1974 | PR3-6 | Hawksnest | A | X | | | | |
| 100-985 | 9114 | 1:20k | 11/14/1977 | PR3-10 | Buck | C | X | | | | X |
| 100-985 | 9115 | 1:20k | 11/14/1977 | PR3-10 | Buck | C | X | | | | X |
| 100-985 | 9116 | 1:20k | 11/14/1977 | PR3-10 | None | X | | | | | |
| 100-985 | 9170 | 1:20k | 11/14/1977 | PR3-10 | Buck | B | X | | | | X |
| 100-985 | 9171 | 1:20k | 11/14/1977 | PR3-10 | Buck | B | | | | | X |
| 100-985 | 9172 | 1:20k | 11/14/1977 | PR3-10 | Buck | B | X | | | | X |
| 100-991 | 5597 | 1:30k | 11/30/1977 | PR3-9 | Buck | B | X | | | | |
| 100-991 | 5598 | 1:30k | 11/30/1977 | PR3-9 | Buck | B | X | | | | |
| 100-991 | 5599 | 1:30k | 11/30/1977 | PR3-9 | Buck | B | X | | | | |
| 100-991 | 5616 | 1:30k | 12/3/1977 | PR3-9 | Buck | B | X | | | | |
| 100-991 | 5617 | 1:30k | 12/3/1977 | PR3-9 | Buck | B | X | | | | |
| 100-993 | 9894 | 1:20k | 12/7/1977 | PR3-10 | Buck | B | X | X | | | |
| 100-993 | 9895 | 1:20k | 12/7/1977 | PR3-10 | Buck | B | | X | | | |
| 100-993 | 9896 | 1:20k | 12/7/1977 | PR3-10 | Buck | B | | X | | | |
| 100-993 | 9897 | 1:20k | 12/7/1977 | PR3-10 | Buck | B | X | X | | | |
| 100-993 | 9898 | 1:20k | 12/7/1977 | PR3-10 | Buck | B | X | | | | |
| 200-221 | 2129 | 1:30k | 3/25/1983 | PR2-3 | Hawksnest | A | | | | | |
| 200-221 | 2129 | 1:30k | 3/25/1983 | PR2-3 | Windswept | A | | | | | |
| 200-221 | 2130 | 1:30k | 3/25/1983 | PR2-3 | Hawksnest | B | | X | | | |
| 200-221 | 2130 | 1:30k | 3/25/1983 | PR2-3 | Windswept | A | | | | | |
| 200-221 | 2137 | 1:30k | 3/25/1983 | PR2-3 | Haulover | A | | | | | |
| 200-221 | 2137 | 1:30k | 3/25/1983 | PR2-3 | Newfound | A | | | | | |
| 200-232 | 2249 | 1:15k | 3/25/1983 | PR2-3 | Newfound | A | | | X | | |

Use of Aerial Photography to Assess Changes in Distribution of Elkhorn Coral in the USVI

| Roll No. | Photo | Scale | Date | Map Index No. | Area Covered | Quality | incomplete | glint | shade | clouds | turbidity |
|----------|-------|-------|-----------|---------------|--------------|---------|------------|-------|-------|--------|-----------|
| 200-232 | 2250 | 1:15k | 3/25/1983 | PR2-3 | Haulover | B | X | | X | | |
| 200-232 | 2250 | 1:15k | 3/25/1983 | PR2-3 | Newfound | A | | | X | | |
| 200-232 | 2251 | 1:15k | 3/25/1983 | PR2-3 | Haulover | B | | | X | | |
| 200-232 | 2251 | 1:15k | 3/25/1983 | PR2-3 | Newfound | A | | | X | | |
| 200-232 | 2285 | 1:15k | 3/25/1983 | PR2-3 | Hawksnest | A | | | X | | |
| 200-232 | 2285 | 1:15k | 3/25/1983 | PR2-3 | Windswept | A | | | | | |
| 200-232 | 2286 | 1:15k | 3/25/1983 | PR2-3 | Hawksnest | A | | | X | | |
| 200-232 | 2286 | 1:15k | 3/25/1983 | PR2-3 | Windswept | A | | | | | |
| 200-569 | 7204 | 1:20k | 1992 | MapFinder | Buck | C | X | | | | |
| 200-569 | 7205 | 1:20k | 1992 | MapFinder | Buck | C | X | | | | |

Figure A: *Acropora palmata* / Study Areas / St. John



Source: 1999 Airphoto mosaic / NOAA



Figure B: *Acropora palmata* Study Area / St. Croix, U.S.V.I.



Buck Island




Source:
NOAA Imagery 1999
Airphoto mosaic

1000 0 1000 Meters


A scale bar with three segments. The first segment is labeled '1000', the second segment is labeled '0', and the third segment is labeled '1000 Meters'.

Figure #1: St. John /Hawksnest Bay / 1974



Legend

 *Acropora palmata*
(60% - 80% cover)

 *Acropora palmata*
($< 10\%$ cover)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application;
- histogram stretch
- brightness & contrast

Date of imagery: 1974
Original scale: 1:12,000




Prepared by
Island Resources Foundation
March 2002

200 0 200 Meters



Figure #2: St. John / Hawksnest /1983



Legend



Acropora palmata
(60% - 80% cover)



Acropora palmata
(< 10% cover)

Note: Interpretation base on
airphoto analysis and Arc View
Image analysis application


Date of imagery: 1983
Original scale: 1:15,000





*Prepared by
Island Resources Foundation
March 2002*



Figure #3: St. John / Hawksnest Bay / 1999



Legend

-  *Acropora palmata* (60% - 80% cover)
-  *Acropora palmata* (<10% cover)

Note: Interpretation based on airphoto analysis and Arc View Image analysis application


Date of imagery: 1999
Original scale: 1:48,000



Prepared by
Island Resources Foundation
March 2002




Figure #4: St. John / Windswept / 1974



Legend



Acropora palmata
(60%-80% cover)



Acropora palmata
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
Image analysis application


NOAA Imagery: 1974
Original scale: 1:12,000




Prepared by
Island Resources Foundation
March 2002




Figure #5: St. John / Windswept / 1983



Legend



Acropora palmata
(60%-80% cover)



Acropora palmata
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
Image analysis application


NOAA Imagery: 1983
Original scale: 1:15,000




Prepared by
Island Resources Foundation
March 2002



Figure #6: St. John / Windswept / 1999



Legend




Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA Imagery: 1999
Original scale: 1:48,000



Prepared by
Island Resources Foundation
April 2002

Figure #7: St. John / Haulover Bay / 1971



Legend

 *Acropora palmata*
(60%- 80% cover)

 *Acropora palmata*
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA imagery: 1971
Original scale: 1:20,000



Prepared by
Island Resources Foundation
April 2002

300 0 300 Meters



Figure #8: St. John / Haulover Bay / 1983



Legend

 *Acropora palmata*
(60% - 80% cover)

 *Acropora palmata*
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA imagery: 1983
Original scale: 1:15,000




Prepared by
Island Resources Foundation
April 2002

300 0 300 Meters



Figure #9: St. John / Haulover Bay / 1999



Legend

 **Acropora palmata**
(**<10% cover**)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA imagery: 1999
Original scale: 1:48,000




Prepared by
Island Resources Foundation
April 2002

300 0 300 Meters



Figure #10: St. John / Newfound Bay / 1971



Legend

 *Acropora palmata*
(60% - 80% cover)

 *Acropora palmata*
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc view
image analysis application

NOAA imagery: 1971
Original scale: 1:20,000




Prepared by
Island Resources Foundation
April 2002

300 0 300 Meters




Figure #11: St. John / Newfound Bay / 1983



Legend

 *Acropora palmata*
(60% - 80% cover)

 *Acropora palmata*
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA imagery: 1983
Original scale: 1:15,000




Prepared by
Island Resources Foundation
April 2002

300 0 300 Meters



Figure #12: St. John / Newfound Bay / 1999



Legend



Acropora palmata
(<10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application


NOAA imagery: 1999
Original scale: 1:48,000



Prepared by
Island Resources Foundation
April 2002



Figure #12a: St. John / Newfound Bay / 1999



Legend

Acropora palmata
low cover (<10%)

Arc view Image analysis
using "level slicing" tool


NOAA imagery: 1999
Original scale: 1:48,000



Prepared by
Island Resources Foundation
April 2002

200 0 200 Meters

Figure #12b: St. John / Newfound Bay / 1999



Legend

Acropora palmata
low cover (<10%)


Arc view Image analysis
using "Gaussian" tool
with higher contrast and
lower brightness

NOAA imagery: 1999
Original scale: 1:48,000



Prepared by
Island Resources Foundation
April 2002

Figure #13: St. Croix / Buck Island / 1977



Legend



Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA imagery: 1977
Original scale: 1:20,000



Prepared by
Island Resources Foundation
May 2002



Figure #14: St. Croix / Buck Island / 1999



Legend

 **Acropora palmata**
(< 10% cover)

Note: Interpretation based on
airphoto analysis and Arc View
image analysis application

NOAA imagery: 1999
Original scale : 1:48,000



Prepared by
Island Resources Foundation
May 2002